

**MATHEMATICS ENRICHMENT CLUB.**  
**Solution Sheet 5, May 30, 2016**

1. By Pythagoras,  $c^2 = a^2 + b^2$ . Hence,  $a^2 = c^2 - b^2 = (c - b)(c + b)$ . So that for  $a^2 = b + c$ , we must have  $b - c = 1$ . Therefore,  $a^2 = b + c = 2b + 1$ , which implies  $a$  is odd because  $b$  is an integer. Let  $k$  be an integer, then  $a$  must be in the form  $a = 2k + 1$ .  
Hence  $a = 2k + 1$ ,  $b = (2k + 1)^2 = 4k^2 + 4k + 1$  and  $c = 4k^2 + 4k + 2$ , for  $k = 1; 2; 3; \dots$  are the solutions.

2. Let  $x = d_0d_1d_2 \dots d_{n-1}d_n$ . Then we can split  $x$  into the sum of two numbers, one consist of the odd digit the other the even digits; That is

$$x = 10^n d_0 + 10^{n-1} d_1 + 10^{n-2} d_2 + \dots + 10 d_{n-1} + d_n$$

$$= (10^n d_0 + 10^{n-2} d_2 + \dots) + (10^{n-1} d_1 + 10^{n-3} d_3 \dots):$$

Now, the remainder of  $10^k$  divided by 11 is 1 when  $k$  is odd, and  $-1$  when  $k$  is even. Recall the properties of remainders [https://en.wikipedia.org/wiki/Modular\\_arithmetic](https://en.wikipedia.org/wiki/Modular_arithmetic). The remainder of  $10^n d_0$  divide by 11 is either  $d_0$  or  $-d_0$  depending on whether  $n$  is even or odd. Therefore, the remainder of  $(10^n d_0 + 10^{n-2} d_2 + \dots)$  divided by 11 is either  $d_0 + d_3 + \dots + d_n$  or  $-(d_0 + d_3 + \dots + d_n)$

- 4.
5. Apply the change of coordinates  $X = x/10$  and  $Y = 10y$ . Then  $X = 10 \cos(10Y)$  and  $Y = 10 \cos 10X$ . In particular, the graph of  $X$  and  $Y$  is symmetric. Let  $A$  be the sum of their  $X$ -coordinate, and  $B$  be the sum of their  $Y$ -coordinate. By the symmetry of graph of  $X$  and  $Y$ , one has  $\frac{A}{B} = 1$ . Moreover, by definition one has  $A = a/10$  and  $B = 10b$ . Hence,  $\frac{a}{b} = \frac{10A}{B/10} = 100$ .
6. Label the points  $p_1; p_2; \dots; p_{100}$ . Draw the  $p_1; \dots; p_{99}$  evenly spaced on a circle in order, and then place the  $p_{100}$  in the center of the circle. Suppose we are able to draw 50 line segments each intersect one another. Then by construction, no lines can pass over more than 2 points. Hence, we may assume without loss of generality, that the points are connect in pairs, and that  $p_1$  is connected to  $p_{100}$ . Consider the point  $p_{50}$ , if it is connected to  $p_k$  for  $1 < k < 50$ , then the line  $p_k p_{50}$  can not possibly intersect  $p_1 p_{100}$  because they belong to different halves of the circle (separated by the diameter pass through  $p_{50}$ ). If  $p_{50}$  is connected to  $p_k$  for  $50 < k < 100$ , then again the lines  $p_k p_{50}$  and  $p_1 p_{100}$  belongs to different halves of the circle (separated by the diameter pass through  $p_{49}$ ).
7. For  $x^x + 1$  to be divisible by  $2^n$ ,  $x^x + 1$  must be even, which implies  $x$  must be odd. Now by using polynomial division argument [https://en.wikipedia.org/wiki/Polynomial\\_long\\_division](https://en.wikipedia.org/wiki/Polynomial_long_division), one can show that

$$x^x + 1 = (x + 1)(x^{x-1} - x^{x-2} + x^{x-3} - \dots + x^2 - x + 1):$$

Since the term  $(x^{x-1} - x^{x-2} + x^{x-3} - \dots + x^2 - x + 1)$  is the sum of odd number of odd numbers, it is an odd number, and therefore can not be divided by  $2^n$ . It follows that  $x + 1$  must be divisible by  $2^n$ ; that is  $x$  must be a multiple of  $2^n - 1$ , so that the least value of  $x$  for which  $x^x + 1$  is divisible by  $2^n$  is  $2^n - 1$ .

### Senior Questions

1. Let the number we are attempting to find be  $n$ . If we add the digits of  $n$ , we get  $1 + 2 + \dots + 8 + 9 = 45$ . Recall that an integer  $n$  is divisible by 9 if and only if the sum of its digits is divisible by 9. Since 9 divides 45, the number  $n$  is always divisible by 9. Thus, our problem is reduce to finding  $n$ , such that  $n$  is divisible by  $99 - 9 = 11$ . Note that we have a divisibility by 11 rule from Q2, so to complete this problem, we just need to arrange the digits of  $n$ , to find the smallest possible combination, such that the sum of the odd digits  $a$  and the sum of the even digits  $b$  of  $n$  satisfies  $a - b = 0 \pmod{11}$ .
2. Since  $a; b; c; d; e$  are consecutive positive integers,  $a = c - 2; b = c - 1; d = c + 1$  and  $e = c + 2$ . So that  $a + b + c + d + e = 5c = x^3$ , for some integer  $x$ . Also,  $b + c + d = 3c = y^2$ , where  $y$  is some positive integer. Since  $5c = x^3$ , and  $c$  is an integer,  $x$  must be a multiple of 5. Hence  $c = 25m^3$  for some integer  $m$ . Now, we know that  $c < 10,000$ ,  $m^3 < 400$ ,  $m \leq 7$ . Since there is only 7 cases for  $m$ , we can easily test them to see which one also satisfies  $3c = y^2$ . The only solution is  $m = 3$ ; that is  $c = 675$ .

3. Let  $y = a + b$ , where  $a = \sqrt[3]{x + \sqrt{x^2 + 1}}$  and  $b = \sqrt[3]{x - \sqrt{x^2 + 1}}$ . Therefore, we wish to find values of  $x$  such that  $y$  is an integer. Note that  $a^3 = x + \sqrt{x^2 + 1}$ ,  $b^3 = x - \sqrt{x^2 + 1}$ . Hence  $a^3 + b^3 = 2x$  and  $a^3 b^3 = -1$ ;  $ab = -1$ . Which implies

$$\begin{aligned} y^3 &= (a + b)^3 \\ &= a^3 + 3a^2b + 3ab^2 + b^3 \\ &= 2x - 3(a + b) \\ &= 2x - 3y \end{aligned}$$

There,  $x = \frac{1}{2}(y^3 + 3y)$  for all integers  $y$ .