Social Policy Research Centre

Research Report

Research on the need for two care workers in a community setting

Final Report

Rosemary Kayess, kylie valentine, Denise Thompson, Ariella Meltzer, Karen R. Fisher

For Ageing, Disability and Home Care,

Department of Family and Community Services NSW

Social Policy Research Centre

May 2013
SPRC Report 5/13

Research team

Social Policy Research Centre, UNSW

Rosemary Kayess, kylie valentine, Denise Thompson, Ariella Meltzer, Karen Fisher

People With Disability Australia

Damien Anderson, Emmie Hallett, Michael Bleasdale

Contact for follow up

Rosemary Kayess, Social Policy Research Centre, University of New South Wales, Sydney NSW 2052, Ph: (02) 9385 7800, r.kayess@unsw.edu.au

Suggested Citation

Two care workers in a community setting

Executive Summary

Ageing, Disability and Home Care (ADHC), Department of Family and Community Services NSW, through its Attendant Care and Physical Disability Unit (ACPDU), commissioned the Social Policy Research Centre (SPRC), University of New South Wales (UNSW) to undertake a research project on the need for two care workers in a community setting.

ADHC wishes to explore approaches to two person services: to understand how services are currently operating, both within the Attendant Care Program (ACP) and the High Needs Pool (HNP), in a community setting, and in hospital and other like settings; to assess the reasons for and the risks of using two workers; to establish the possible use of equipment and technological solutions and their cost; and to develop appropriate training packages and guidelines.

The methodology was a review of literature and policies and consultations with interested stakeholders, the development of guidelines and a request form template, and a report. This report is a summary of the findings. The guidelines are a separate document with specific considerations for determining the need and alternatives to two care workers.

Literature review

The literature review was about when are two or more attendant care workers needed to work at the same task at the same time in a community setting. The usual context within which the use of two person care is mentioned involves managing lifting, moving and transferring clients, risk assessment and safe handling. This section summarises the literature, with the full references included in the body of the report.

There is very little mention of two person care in the community care literature, and nothing at all on two person care (or care by more than one person) as a separate issue. When the presence of more than one worker is mentioned, it is not distinguished from one person care. The review did not uncover any instances where care by more than one person was mandated across the board in specified circumstances.

Most of the literature about safe client-handling and risk of staff injur

There is some discrepancy among the findings of various studies about the extent to which team lifting can reduce musculoskeletal injuries among care workers. A number of studies have shown that that it does, but at least one study failed to find some reduction in employee injuries after the introduction of team lifting. Other research has also found that having two or more workers is no guarantee of avoiding injury. The guidelines for avoiding musculoskeletal injuries among care workers do not indicate that injuries only occur when staff work alone.

The current emphasis in the human services sector is on eliminating client lifting to the fullest extent possible, whether performed by one person or by more than one. The usual term for this is ‘No Lifting’, and it involves:

- encouraging clients to assist in their own transfers
- assessments of client-handling risk
- using mechanical lifting aids and other equipment
- modifying the work environment to accommodate the equipment and
- training all staff in the correct use of equipment and techniques for moving and transferring clients.

Stakeholder views

The second research method was interviews with interested stakeholders, including clients, family members, service providers, advocacy and service peaks and government representatives. Most stakeholders noted that the use of two care workers has become more common. This was attributed to people with disabilities increasingly living within the community resulting in an increased complexity of support needs of people living at home as well as an increased emphasis on workplace safety. The most frequent reasons for the use of two care workers were:

- Client age, weight, size, and physical function (capacity to assist with lifting and positioning)
- Client challenging behaviour
- Scoliosis, spasticity or other high support needs related to positioning.

The stakeholders described the primary benefits of two care workers as relating to worker and client safety. They thought that using two care workers reduces the risk of manual handling injuries and unsafe handling of clients, and provides a more secure working environment for workers, in which the risks to workers from clients’ challenging behaviour, grievances and complaints, or unsafe environments are lessened. The perception of safety seems to be as important to workers and clients as actual risk. The interviews indicate they do not agree about whether the use of two care workers actually reduces the risk of injuries, compared to the use of one worker.

They identified assessment as an area in need of improvement. In terms of indicators for the use of two care workers, stakeholders identified the need for risk assessments to include:

and solutions, and address both the actual and perceived benefits and risks of two care workers.

The principles to make decisions about alternatives to two care workers relate to:

- client and worker safety
- meeting the needs of clients and
- ensuring that both clients and workers feel secure.

Some of the alternatives to two care workers relate to training and equipment. Other alternatives relate to better systems of support for workers and clients, to address the anxieties about the use of single

Two care workers in a community setting

Two care workers in a community setting

2.3 Steering committee

Another source of data was the project's steering committee, which was constituted to provide advice and feedback on the project's methodology and findings. The steering committee was made up of representatives of consumer, policy and service delivery organisations: Northwest Therapy, ADHC Manual Handling Unit, Brain Injury Association of NSW, Attendant Care Industry Association, Physical Disability Council of NSW, Spinal Cord Injuries Association, Just Better Care Community Services, Australian Home Care Service, Lifetime Care and Support Authority, MS Australia, Cerebral Palsy Alliance.

2.4 Limitations and caveats

The sample for the consultations was very small, and participated after nomination from ADHC. The views of participants cannot be generalised, as it is not possible to know how representative their experiences and opinions are.

In particular, there are limitations to the client sample in terms of age, as indicated above, and limitations in that most clients (except one) had a physical disability. Given the issue of challenging behaviour reflected in this report, not including other disability types is seen as a limitation.

Further, there are limitations to the care worker sample. Two care workers were interviewed in this sample. This number did not allow full reflection of a range of issues for and characteristics of care workers that may affect their perspective on the two care worker issue, such as their own size and strength, type of training, type of agency, experiences in service provision and philosophy of service or care.
Future J 0 Tc B2(he(r)7(ai)1507(e.)9s24(d)10(ddpr)7(es)4ns)4( )10(t)2h Tc B2(a l)6(i)6(m)-3(i)6(t)1

3 Literature and policy review

3.1 Definitions

The expansion of the research to when are two or more attendant care workers needed to work at the same task at the same time in a community setting focused on identifying the occasions on which any increase in service hours is justified because there is no viable alternative.

The inclusion of the term 'two or more workers' is a response to the discussion in the literature about OH&S issues in the healthcare workforce that occurs in the context of institutional care – hospitals, aged care and long-term care facilities – where there are many members of staff present all the time. In that context, questions about whether tasks should be performed by one or more people make reference to more than two staff members.

Mention of the fact that the two (or more) employees are working 'at the same task' and 'at the same time' involves a further clarification. Simply referring to 'two (or more) workers' is not sufficiently precise. Clients are often served by more than one worker. For example, ADHC's guidelines for manual handling risk management in community care (DADHC, 2006: 18) notes that a client could have up to 10 staff working in their home, counting night shift, weekend staff and casual staff as well as the client's usual worker(s). However, these staff do not all work at the same time.

The attendant care worker is usually the sole worker providing assistance at any one time for the person with disability. For example, the fact that only one staff member is to be provided per service is emphasised in the assessment procedures of Personal Attendant Care Inc (PAC, 2009b), a charitable non-profit organisation in Ontario, Canada; and the wording of their health and safety regulations in relation to lifts and transfers (PAC, 2009a) makes it quite clear that services are being carried out by a single worker.

Working singly is recognised as one of the organisational aspects of the job that can increase the level of risk. In their Health and Safety Guidelines for Home Care Workers(UNISON, no date), Britain's largest public sector union commented that 'As lone workers, home care workers can be particularly vulnerable as they are isolated from other workers and cannot easily liaise with colleagues'. Again, the Queensland Government's Code of Practice for manual tasks involving the handling of people (Queensland Government, 2001) acknowledges that working singly means that a worker cannot call on another person for assistance or use team-handling. However, there is no suggestion that the risks should be managed by supplying two workers where the task has traditionally been done by one.

It may sometimes be necessary, of course, for two people to be available for the same task at the same time, and a number of examples are given in what follows. However, each of these examples involves a particular situation and there is no suggestion that the use of two workers should be made into a mandatory requirement across the board, e.g. whenever there is a hoist involved, without the need to consider each case on its merits.

with nursing staff in hospitals, even though work in private homes entails many of the same risks as institutional health care, as well as extra burdens because the work environment is uncontrolled and less standardised (Ono et al, 1995). (There is some Swedish research on ergonomic risks to home care workers – see section 2.4 below).

The review did not uncover any instances where two person care was mandated in specified circumstances across the board, for example, when hoists were being used. The Victorian NW Residential Support Services guidelines on manual assistance to people in residential care (2001), for example, do not even mention two person care. Instead, recommendations refer to the need for assessing each particular case as it presented. A manual-handling care plan for each person is the recommended procedure. These guidelines state that the plan should indicate the specific manual-assistance techniques the support worker should use for particular activities. They also specifically forbid the use of particular lifting techniques (except in an emergency), i.e. shoulder lift, cradle lift, top-and-tail lift, and lifting a resident up from the floor on one's own. But they do not specify any situations where two support workers should be used (Victorian NW Residential Support Services, 2001). Similarly, although the UK Department of Health's national minimum standards specify that 'two people fully trained in current safe handling techniques and the equipment to be used are … involved in the provision of care', this is only the case when the need has been identified from the manual-handling risk assessment(UK DoH, 2003).

3.3 Institutional setting

Most of the literature on safe client handling, and on the risks of injuries to staff, refers to institutional care – hospitals, aged care facilities – rather than to community settings, e.g. the O'Shea No Lift System (see below), O'Shea and Hennessey's Handbook for WorkSafe Victoria, Transferring People Safely (O'Shea and Hennessey, 2009), which was produced specifically for use by Victorian employers in health and aged care settings.

Legislation has not been clear who is responsible for the condition of the premises when the private home becomes a workplace as in the case of attendant care. Most service providers hold clients responsible for maintaining safe premises, and include that in their service agreements as a condition for the provision of services. However, a better approach is for services providers, workers and clients to work in partnership to consider the risks and how to best address them, and a formal framework to guide collaboration is useful (DADHC, 2006).

In NSW (and elsewhere) the key features of a client-handling policy are:

- that workers are not to manually move a client's body weight, either wholly or in part, unaided
- that mechanical equipment must be used if a client has to be moved and
- that manual lifting is to be used only in extreme circumstances, either in emergencies or when all other methods have failed (DADHC, 2006).

The NSW government, through ADHC, has developed a manual handling policy for disability and community care supported by Best Practice Guidelines for manual-handling risk management (DADHC, 2006). There was a review of the manual-handling training programs of the Home Care Service of NSW, a statutory corporation and the largest provider of home-based care in NSW. In January 2005, ADHC introduced a program using occupational therapy manual-handling advisors to conduct expert risk assessments with a view to eliminating high-risk tasks from staff routines. The program targeted sites with high exposure to manual handling risk and a history of manual-handling incidents and injuries. Manual-handling injuries and the costs associated with them have been reduced since 1997/98(WorkCover NSW, 2006).

There is general agreement at the policy level that team lifting as a way of dealing with manual handling risks should be used only as a last resort. The NSW Occupational Health and Safety Regulation 2001, for example, emphasises the importance of achieving risk control by means other than team lifting, as far as is reasonably practicable. ADHC's 'DisabilitySafe' website[4] stresses the same point, suggesting a number of other ways of controlling manual handling risks, e.g. by modifying the work environment or the design of objects, by providing mechanical aids, by ensuring staff are adequately trained. (See also: WorkCover NSW, 2004; WorkCover NSW, 2006).

The Queensland Government's Code of Practice for manual handling tasks (Queensland Government, 2001) also says that team handling should only be used when there is no other solution. The authors go further to say that team handling itself involves a number of risks, namely:

- different capacities and physical dimensions of team members
- different degrees or timing of force exerted by different team members
- loss of coordination due to adjustments made to fit in with other team members
-

The Queensland Government's Code of Practice for the handling of people (Queensland Government, 2001) also refers to using more than one worker in the short term while waiting for a better solution. For example, when a device is not immediately available, team-handling with training might be used until the mechanical device is acquired. As well, the Code referred to two or more workers in other contexts:

- during a sling lift from the floor with one worker operating the sling while the other worker communicates with and reassures the person and assists where necessary

- in the case of individual characteristics of the worker in a team handling situation – younger, older, pregnant, with an existing back injury and

- raising a person from the floor in an aged care residential facility.

The US Occupational Safety and Health Administration's Guidelines for Nursing Homes (OSHA, 2009) says that the number of carers required depends in part on the size and weight of the person being moved(as well as which equipment is needed). The Guidelines describe a number of occasions when more than one carer is needed:

- when clients cannot help to reposition themselves in a Geri or Cardiac chair (a friction-reducing device is also needed)

- when using a ceiling hoist (although the Guidelines also note that some residents can use this kind of device without assistance)

- in the case of lateral transfers or

)11.82d [(i)6(a)10(4(f1-8(er)7(s)4()bod))as)4(d(ar)7(n a c)bTJd4

two (or more) carers that was most frequently mentioned was the client's uncooperativeness; other factors were weight (i.e. >200 pounds), the inability to assist, and complicated clinical conditions.

Care of bariatric clients

One type of situation which is most likely to require two or more workers is the care of bariatric clients in institutional settings.[8] Moving clients who are large, (bariatric clients often classified as those more than 30% over Body Mass Index [BMI] with a medical problem) whether by weight or height, poses even higher risks than moving people of average height and weight. The South Australian government's Guidelines for community workers (SA Government, 2009) advises making sure that there enough workers to move bariatric clients. The Disability Safe website[9] emphasises the importance of including clients' sizes as part of the assessment procedure and of ensuring that the equipment available matches the tasks which need to be done. They also recommend consulting the suppliers of equipment specifically for bariatric clients when developing manual handling plans for such clients.

Once again, the focus of most of the literature is on institutional settings (e.g. Weinel, 2008), although much of what is discussed is relevant to the management of bariatric clients in their own homes. Muir and Archer-Heese (2009), for example, point out that even raising a client's limb for a dressing change can be dangerous. The weight of a leg of a 350-pound client, they say, would be 62 pounds, well over the recommended maximum of 35 pounds. In this case, staff should use a mechanical lift device and a limb sling. They recommend a ceiling lift as the best choice, both for bariatric transfers and repositioning someone in bed, and for reducing space requirements. They point out that the key to effective and safe bariatric-client handling is thorough preparation prior to admission (or in the case of attendant care, as part of the assessment process). Following the safe bariatric patient-handling toolkit produced by the US Department of Veterans' Affairs,[10] they list a number of essential components of a safe program, namely:

- operational procedure and policy
- patient assessment tools
- communication tools
- patient handling algorithms and guidelines
- space and environment considerations
- equipment needs

---

[8] Bariatrics is the science of providing healthcare for those who have extreme obesity (Muir and Archer-Heese, 2009).

[9] http://www.disabilitysafe.com.au/hazards-risks/manual-handling

[10] http://www.visn8.va.gov/visn8/patientsafetycenter/safePtHandling/toolkitBariatrics.asp

- staff training and education and
- evaluation.

These authors do not discuss the need for two or more care workers in this context, apart from mentioning that using an older floor lift requires two staff to move it safely in transferring a bariatric client from bed to chair, and a third staff member to monitor the client (Muir and Archer-Heese, 2009).

The founder and president of American Bariatric Consultants,[11]Kevin Huffman (Carlson, 2008), is also conscious of the fact that arrangements need to be made to care for bariatric clients in their own homes. He advises occupational therapists to try and ensure that their bariatric clients who are beariTel 0.038 T3.21(517 0 Td [(to)-1w)16(nnc

for all settings; they require infrastructure and equipment to support their use; and staffing needs careful consideration. However, most of these issues related to an institutional setting rather than community care.

A systematic review of research reports on intervention strategies to reduce the risk

transferring a client from a bed to a chair, given a choice between solo lift, two person lift, lifting team, mechanical lift and other. Musculoskeletal problems were identified as pain, numbness, tingling, aching, stiffness, or burning in the neck, shoulder and back. The study found that all the preventive devices – lifting teams, mechanical lifting devices, adjustable beds, and sliding/transfer boards – were significantly related to musculoskeletal disorders of the back, although lower levels of disorders were associated with lifting teams and lifting devices than with the other means of client transfer. Transfer boards/sliding sheets and adjustable beds were associated with significantly higher levels of back disorders. In the case of neck disorders, only the lifting devices made any significant difference in lowering the risk of injury. The transfer method most popular with the hospital nurses was the two person lift, although it was less popular with nurses in nursing homes and home health agencies, who preferred mechanical lifts. Those who had been trained in the use of lifting devices were twice as likely as the rest of the sample to prefer using these devices, an outcome which the authors referred to as ‘encouraging’.

Other US studies have also found that the use of lift teams leads to cost savings due to reductions in lost work time and in claims for compensation. One study (Charney et al, 1991) found, for example, that injury rates dropped from 39 per 1000 to 2.4 per 1000 in one year at a US hospital after the introduction of lift teams, thus saving $65,000 on the day shift; another study (Charney, 1997) found that accidents resulting in lost time dropped from 16 to one in another large hospital, thus saving $144,000 in the year following the introduction of teams.

In contrast to all the studies showing that the introduction of team lifting reduced injuries and musculoskeletal disorders among nursing staff, another US study (Springer et al, 2009) failed to find same reduction in employee injuries described by previous authors. The authors of this study explained the discrepancy in terms of the length of time over which the comparisons were made. They compared the level oC 0.010((c)4(s)4(i)6(ng)10)6(ngm)-3(s)14(.)]T 30.14ID 2 >>BDC 10( )]11.487(s)]TJ or

while the two carer procedure is rated ‘F’). The other manual lifting tasks assessed in the guide as high risk are not identified by the number of people performing them. It might be assumed, then, that they are likely to cause injury, and hence are not recommended, however many carers perform them. In other words, the criterion of risk reduction in any task is not the number of people who perform it, but the availability of mechanical aids (slide sheets in the case of turning or re-positioning in bed). The guide did refer to a number of procedures requiring more than one person to perform:

- moving a client f 0.(g)10l 6or11pt 0uou0(dnTJ(en)]TJ)Tje12 -2.22 Tdu6 supp6(n2(n b)10(d

However, although manual handling involves more than lifting – it involves pushing, pulling, lowering, holding, carrying and restraining as well as lifting (WorkCover NSW, 2006) –'No Lifting' still appears to be the preferred terminology.

The Victorian Department of Health (2004), for example, uses the term 'No Lifting' and describes it as the elimination of the manual lifting of clients in all but exceptional or life-

nurses to be proactive in identifying hazards and reducing risks of injury in the workplace (Keating et al, 2002).

community settings and private homes, as well as in acute hospitals and aged care facilities.

It would appear that there has been increased interest in ceiling hoists as part of the safe handling approach to decreasing the physical exertion of carers (Jung and Bridge, 2009). WorkSafe Victoria (2009) prefers ceiling hoists ('overhead track systems') for moving clients, noting that ceiling hoists require staff to exert significantly less force than mobile, floor-based hoists. The overhead hoists are more efficient, take less time to transfer a person, and they are more acceptable to clients. In the UK, Hall (2002) pointed out that an overhead hoist usually just required the worker to operate a handset and guide the carry-bar into place, rather than having to support the weight of either the client or the carry-bar. (Only one worker was mentioned in this context). WorkSafe Victoria (2009) recommended that ceiling hoists be installed in all new and renovated health and aged care facilities where people needed to be transferred, but private homes were not mentioned.

Writing in the context of an evaluation of the use of ceiling hoists in a newly built spinal cord injury unit at a veterans' hospital in Florida, Weinel (2008) noted that ceiling-mounted lifts were a viable alternative to floor-based lifts. The lifts use tracks mounted on overhead beams sturdy enough to support someone's weight. They have battery-powered lifting motors which raise and lower the clients and move them along the tracks, which can be either single or H-shaped so as the cover a wider area. Some of the motor units are portable so that they can be relocated from one track to another in another room. The client is suspended from the motor unit in a sling and there is a variety of sling designs and fabrics available. The advantage of a ceiling lift where frequent lifts and transfers are necessary is that it is always accessible. The nursing staff in this facility preferred a two-function (up-down) control rather than the multi-function powered tracking. They felt that the powered tracking motor was too slow and they favoured a hands-on approach, especially as they could move the person in the sling along the track with very little effort. Clients reported feeling secure during transfer and being less jostled than with the floor lift.

The author (Weinel, 2008) said that studies comparing floor lifts with ceiling lifts had found that there were fewer musculoskeletal injuries among staff with use of the ceiling lift. Other studies had found that ceiling lifts required half the effort of floor lifts, and that lifting and transferring with a ceiling lift produced less trunk and shoulder muscle activity than with a floor lift.

The Victorian Hospitals Industrial Association (VHIA, 2003) listed the key components of what they called 'overhead tracking' for an institutional setting as: a ceiling track; an electric motor; a suspended sling or frame; and handset control. The authors said that there were a number of different forms of tracking, namely:

- straight or curved, and in various lengths
- with turntable junctions for changes in direction
- a traverse or H-shaped system which allows for movements both up/down and across

point out that significant numbers of incidents are reported each year where people have been injured while being moved with hoisting equipment. It lists a number of things that can go wrong, namely:

- the wrong size sling – discomfort if it is too small and the risk of the person slipping out if it is too large
- the wrong type of hoist or sling for the particular person or the specific task – inadequate support and increased risk of falling
- incompatibility of the hoist and sling – insecure attachment between the two
- failure of equipment due to poor maintenance, lack of inspection, inappropriate laundering, or inadequate repair or modification
- leaving someone unattended in a hoist
- hoist overturning on a difficult surface and
- not using the safety harness/attachment (UK HSE, 2011).

The information sheet provides a checklist for safe handling practice of 24 items – e.g. 'Review the person's condition prior to each transfer to ensure continued suitability' – together with additional guidance separately for mobile hoists, ceiling track/overhead systems, standing hoists, slings and bath hoists. It also provides a checklist for equipment safety checks prior to each use.

As already mentioned, there is little literature on the use of ceiling hoists in private homes. The authors of the above-mentioned review (Jung and Bridge, 2009) said that ceiling hoists were given low priority in community care strategies, even though it was an important home modification issue. They surmised that ceiling hoists were so rarely considered for people being cared for at home, because the systems were initially introduced in response to the high incidence of back injuries among nurses.

great deal to say about them. The installation of a ceiling hoist was mentioned only on two occasions. It was one among a wide range of house modifications mentioned by focus group participants; and one participant reported that the owners of the house he occupied, the public housing authority Housing New Zealand, had refused to allow a ceiling hoist to be installed, even though he had been assessed as needing it because the house was not suitable for a wheelchair. The research was commissioned by the Centre for Housing Research, Aotearoa New Zealand and carried out by the Auckland Disability Resource Centre. The report (Saville-Smith et al, 2007) mentioned ceiling hoists in the context of a wider concern with ways of optimising housing access for people with moderate to severe physical and sensory disabilities affecting their mobility. The research focused on: the current housing experiences of people with physical disabilities; the capacity of the housing market to respond to the demand for 'lifetime homes'; and the opportunities to establish a housing stock that was 'future-proofed' for those with moderate to severe physical disabilities. The chief finding of the research was that the housing needs of people with disabilities were not being met by New Zealand's current housing stock.

As already noted, the issue of the need for two (or more) care workers in an attendant care setting is not discussed at any length in the literature. However, there are a few fleeting mentions. The VHIA (2003) said that using-

3.10 Assessment

The importance of assessment is stressed throughout the literature. As the authors of the first evaluation of the VNBIPP (Keating et al, 2002) said, assessment of client-handling requirements is a prerequisite for identifying and reducing the risks associated with client handling.

There are a number of sources which specify what should be included in any process of assessment for client-handling risks. The report of the second VNBIPP evaluation (Martin et al, 2004) stressed the importance of identifying the needs and abilities of clients when assessing requirements for lifting and repositioning tasks.

WorkCover NSW (2006) points out that risk assessments in the health industry are more complex than in other workplaces. In most work environments the only assessment needed relates to inanimate objects, which do not change their size, shape or weight from one occasion to another. However, people are not rigid or stable, they are not always the same shape every time they are handled, and consideration must also be given to attributes such as physical ability, mental status and cognition, medical condition and communication issues. WorkCover NSW (2006) lists four main areas of risk assessment required for handling people:

- the manual-handling task
- the patient-handling risk
- the workplace including the home and
- the equipment.

People also need to be moved in many different ways:

- moving up and down and repositioning in bed
- moving in and out of bed or trolley
- getting into and out of a chair
- moving to and from toilet or commode chair and
- walking.

The risk involved in each of these tasks needs to be assessed, and the assessment repeated at regular intervals, depending on changes in the person's condition. The assessment should be undertaken by the people who are doing the work or at least in consultation with them.

The South Australian government's guidelines (SA government, 2009) listed the following factors to be considered when identifying manual task hazards for community workers:

- actions, postures and movements e.g. bending, twisting, over-stretching

- workplace conditions, e.g. cramped work space, low work surface, uneven or slippery floor surfaces, poor lighting, extremes of hot or cold
- weights and forces (worker should not lift more than 16-20kg)
- characteristics of the load, e.g. unstable or unpredictable, difficult to slide, push, pull or turn
- location of load and distances moved, e.g. storage above shoulder or below knee or load carried a long distance
- frequent and prolonged movements e.g. repetitive tasks, prolonged exertion
- job organisation, e.g. heavy workload, too many clients in one day, lack of staff, unrealistic deadlines, bottlenecks of work and
- individual factors, e.g. worker skills and training, worker hampered by illness, disability or restrictive clothing.

The US OSHA (2009) lists for inclusion in the client assessment:

- the level of assistance required
- the person's size and weight
- the person's ability and willingness to understand and cooperate and
- any medical conditions that may influence the choice of methods for lifting or repositioning.

In the UK, UNISON (no date) advises employers to consider the following in assessing manual-handling risks to home care workers:

- floor surfaces, e.g. uneven, slippery, include steps
- storage arrangements
- the size and layout of the client's home
- the type and size of equipmentha(e)10(c)ehe UK i 222isa8>Tj /TT0 1 Tf0 Td ( )Tj EMC /F.0

evaluation of the environment and of the handling aids required, as well as a client care-plan containing clear information on the client's movement abilities and needs, instructions on handling techniques and aids, and the number of nurses to be used for various moves.

The second level of assessment listed by the guide was an assessment of the hospital department or ward. This does not translate directly into attendant care because every client's home is different, unlike the standardised layouts of hospital wards. Nonetheless, there is still a range of information that can be gathered at this level, from space in the bathroom and the availability of handling aids to the training of staff. The guide includes an example of a form designed to collect information for such an assessment.

The third level of assessment, the top level, referred to senior management considering the requirements of the organisation as a whole, e.g. decisions on budgets, training, uniforms. For hospitals, much of the necessary information is collected as a matter of course, but for care in the community, a survey might need to be done to evaluate general needs.

The guide (RCN, 2003) noted that attendant care providers could minimise the amount of local assessment required for the second level of assessment by producing generic assessments wherever possible. These could also serve as guides for attendant care workers to use in assessing particular manoeuvres in people's homes. Possible areas for generic assessments are:

- transfers from and to bed, chair, commode, toilet
- working with clients who have a history of falls
- bathing
- the condition of floors (hoists on carpets, slippery bathroom floors)
- difficulties in using hoists, e.g. carpets, restricted space
- transfers into and out of car
- babies in high-sided cots and
- handling supplies (packages/boxes) at health centres.

In the UK, the Department of Health's National Minimum Standards for domiciliary care (i.e. attendant care) (UK DoH, 2003) states that a detailed assessment of the risks associated with the delivery of the service must be carried out by the service provider for each new home. The registered person[14] ensures that the assessment is undertaken, by a trained and qualified person, before the attendant care worker

[14] 'All agencies providing personal domiciliary care services, irrespective of size will be required to have a person as registered as the "Fit Person" who has overall responsibility for the service. This person may be the owner or the most senior manager of the service' (UK DoH, 2003: 5).

commences work, and that it is updated annually or more frequently if necessary. A plan for managing the risks is drawn up in consultation with the service user and their family or representatives. It is included in the overall service-user plan and kept in the home of the service user for staff to refer to. Assessment of the risks associated with manual handling must be undertaken separately. The only mention of a need for two person care occurs in this context of assessment: ‘Two people fully trained in current safe handling techniques and the equipment to be used are always involved in the provision of care when the need is identified from the manual handling risk assessment’(UK DoH, 2003: 24, Standard 12.8).

There are many manual-handling risk assessment tools in the literature.

- client-handling equipment and devices
- policy initiatives, the three primary examples being ‘no lift’ policies, ergonomic assessments of client care areas, and the use of client lift teams
- training of staff in the safe use of patient handling equipment, to the use of peer leaders, and to clinical tools such as patient-assessment protocols and algorithms.

An algorithm is a sequence of logical steps to be taken to achieve a given task, consisting of a finite list of precisely defined successive ‘if ... then’ decision points displayed in a flow chart. For example, in the case of the task of transferring someone to or from bed, chair, toilet or car, the algorithm would start with the question, ‘Can client assist?’ If the answer is ‘yes’, no carer assistance is required apart from standing by for safety as needed. If the answer is ‘no’, either wholly or in part, the next decision point relates to the client’s weight. If the client weighs more than 200 pounds the specified action is to use a friction reducing device plus three carers; if less, the action is only to use a friction reducing device. (For more detail about algorithms, see: Fragala et al, 2001; Nelson et al, 2003; Nelson, ed., 2006; and for a list of examples see Appendix 1).

A trial of a ‘best practices’ musculoskeletal injury-prevention program in six nursing homes over six years (Collins et al, 2004) found that the program had had a number of positive results. The program involved the introduction of mechanical lifts and repositioning aids, a zero-lift policy, and employee training in lift usage. After the program, the rate, severity and cost of injuries associated with lifting and moving residents, lost and restricted work days, and repeated staff injuries, were all significantly reduced in all the nursing homes, for staff in all age groups and lengths and types of tenure. Assaults and violent acts by residents towards staff also declined. The reduction in workers’ compensation claims recovered the initial capital outlay in slightly under three years.

A systematic review of studies investigating intervention strategies to reduce the risks associated with patient handling activities (Hignett, 2003) identified the seven most commonly used strategies:

- equipment provision
- education and training in, for example, risk assessment, use of equipment and client assessment
- the introduction of risk assessment
- the introduction of safe-handling policies and procedures
- the introduction of a client assessment system
- redesigning of the work environment and
- changes in work organisation and practices.

Two care workers in a community setting

disability to live and participate in their family and community. The program policy is directed at increasing the independence of the client, and, or reducing risk of illness or injury to either the client or attendant care worker.

The EnableNSW Aids and Equipment program is means tested and operates on a consumer copayment system: a contribution of a fixed $100 per year for pension recipients, low income adults and children under 16 years or for those in a higher income band 20% of the cost. EnableNSW set no upper limits on the cost of individual assistive technology equipment but the item must be the most basic that meets the person's need.

Equipment can be available for loan through the EnableNSW equipment loan pool for trial. The program is managed locally through Local Health Networks with services available local area health services. The assessment and trial can also be facilitated through suppliers making available on loan or through rental pieces of equipment prior to purchase. The ongoing maintenance of equipment is generally available through the supplier and required either on a 6 monthly or 12 monthly basis. Some equipment may come with a warranty that covers maintenance costs for an initial period, say 3 or 5 yrs. Others may require service calls which would include a service call charge and any parts or labour. These cost outlays are very dependent upon the particular piece of equipment and the supplier. Equipment that is sourced through EnableNSW Aids and Equipment program maintenance is covered by the scheme.

Cost implications of equipment include the assessment/trial, purchase and ongoing maintenance of capital equipment costs (Table 3.1).

Table 3.1: Cost implications for the trial, purchase and maintenance of equipment

Purchase Cost	Example	Trial	Maintenance

4 Stakeholder consultation

This section summarises the views of the stakeholders about the research questions. The final section draws on these findings to discuss implications for

The complexity of interpersonal relationships is mirrored in the complexity of organisational factors with two workers. This relates to staff recruitment, rosters and leave. Seven stakeholders from peak bodies, government and service provision agencies talked about the difficulties of organising two workers rather than one, and the difficulties that arise when one worker is sick or arrives late, or when workers do not get on and cannot work well together. While cost is obviously a significant factor in terms of salaries, the consultations suggest that the indirect costs of two carers, associated with the organisational resources required to coordinate this arrangement, are also significant.

4.5 Alternatives to two care workers

There were no alternatives to the use of two attendant carers proposed for clients who use ventilators. A meeting of the reference group noted that the literature describes glossopharyngeal breathing, or ‘frog breathing’ as a technique used by some people who are otherwise fully ventilator dependent (Maltais, 2011). In circumstances where a client does not want or require two attendant care workers except as a safety measure around ventilation, and has been trained in this technique, two attendant care workers may not be needed. However, the consultations and literature suggest that this circumstance is rare.

The stakeholder consultations identified possible alternatives to two attendant care workers for all clients, with the exception of those who use ventiltors. When the decision to use two care workers is based on physical considerations—the nature of the client’s physical disability, and so the nature of the physical tasks undertaken by workers—the alternatives proposed were equipment, environmental modifications, and staff training. Challenging behaviour and worker safety also recurred as a reason for two workers, and fewer alternatives were suggested here,

Reassessment of equipment at intervals was also recommended. For example, clients who received an injury twenty years ago may be using outdated equipment.

> includes all of those parameters, including the skill level of the staff involved, and the training and those sorts of things as well. There isn't a blanket, 'You need two people to use a hoist', no, no.

Representatives of government agencies also emphasised the utility of training about effective use of equipment such as hoists, so that one worker is sufficient in at least some circumstances.

Training does not relate only to use of equipment, but also to the physical training of care workers, for example education in lifting techniques, warm-up exercises and stretching. It also relates to training in specific contexts, particularly manual handling of clients in domestic rather than institutional environments. Workers familiar with hospital settings, or who received training that assumed their workplaces would be group homes or hospitals, need specific, problem-solving based training in home environments. This may involve the assessment of the home's potential for modification, including furniture, floor coverings and load-bearing capacity of walls and ceilings. It may also involve trialling the use of different kinds of equipment. Support and training is necessary in order for workers to make these specific, responsive assessments to individual clients and environments.

Challenging behaviour

Challenging behaviour is defined as behaviour of 'such intensity, frequency or duration that the physical safety of the person or others is placed in serious jeopardy or behaviour which is likely to seriously limit or deny access to the use of ordinary community facilities' (Emerson, 1995). Half the respondents raised client challenging behaviour , as one of the reasons for two care workers (10/23 interviews), although the research literature about this is sparse. This suggests a gap in the research literature, perhaps because challenging behaviour is becoming more concerning to workers and agencies, and research has yet to catch up with this. In circumstances where the worker is at risk of being subject to the effects of a client's challenging behaviour, intervention programs for the client were suggested by one stakeholder. The research literature indicates that interventions based on functional assessments of challenging behaviour and positive behaviour support

as team management structures to provide peer support for validation and learning can address isolation and confidence of workers. Such strategies are also opportunities to follow the principles of person centred planning, by building a team in which the client's specific needs are at the centre, and in which communication, training and support for staff are based on these needs.

The concerns expressed by clients relate in some cases to safety, especially at times of transition, such as moving from an institutional environment, where teams of workers are common, to a home environment. Identifying and responding to client anxiety around the use of single workers is also a positive step. In other cases, cultural and social concerns may be responsible for client anxiety: for example, a strong preference by some clients for female or male care worker, or the fear that some clients have of being alone with a single worker. Again, the responses to these concerns can be guided by the principles of solving specific problems and responding to individual needs: for example, traumatic responses to previous abuse require more nuanced responses than resorting to two care workers.

Risk management is an important consideration for agencies, particularly in relation to the risk of client and worker injury. Addressing the responsibilities of agencies to minimise risk, and adapting the sometimes inflexible systems in which risk assessments are devised, can avoid a simplistic two care worker response to risk management.

Table 5.1 summarises the principles and strategies discussed in this section of the report. It lists the principles behind decisions about the use of two care workers, as identified in the literature and stakeholder consultation. It also identifies strategies to consider the feasibility of 4(i)6(d)10(t)2(r2a as)4( )]TJ eion(k)-64(on(k)-64(on(k)-64(on4)-10u)6

Table 5.1: Principles and alternative strategies instead of two care workers for consideration of the feasibility in the particular circumstances of the person

Reason for two care workers	Principle behind the reason	Alternative strategies for consideration of the feasibility to address the reason and principle in the particular circumstances of the person
Client size, weight, disability	Client safety	Assessment process: tasks required, level of client physical functioning, home environment, skills/techniques/equipment needed
		Home modification to install equipment
		Trial use of equipment (via loan pool, assistance from equipment agency representative)
		Installation of high quality equipment e.g. hoists, shower commodes, trolleys
		Training in the use of equipment
		In situ training in home environments
	Worker safety	Physical matching of client and worker
		Two workers as an interim measure to determine worker characteristics (strength, flexibility, height) needed
	Quality of work	Peer support networks
		Opportunities for feedback and validation
		Regular monitoring and retraining opportunities
Challenging behaviour	Worker safety	Staff training to ensure a consistent approach
		Functional assessment (what functions does the behaviour serve for the client? What is being communicated by the behaviour?)

Maier, J. M. (no date) ‘Proper client transfer technique’ eHow Health– http://www.ehow.com/way_5655339_proper-client-transfer-technique.html

Maltais, F. (2011), 'Glossopharyngeal Breathing', American Journal of Respiratory and Critical Care Medicine, 184(3), August 1, 2011, 381.

Marras, W. S., K. G. Davis, B. C. Kirking and P. K. Bertsche (1999) ‘A comprehensive analysis of low-back disorder risk and spinal loading during the transferring and repositioning of clients using different techniques’ Ergonomics 42(7): 904-26

Martin, P. J., J. T. Harvey, J. F. Culvenor, W. R. Payne and D. Else (2004) VictorianNursesBackInjuryPreventionProjectevaluation report. Melbourne: Policy and Strategic Projects Division, Victorian Government Department of Human Services, Melbourne, Victoria – http://www.health.vic.gov.au/nursing/promoting/back_injury

Matz, M. (no date) ‘Client handling equipment coverage and space recommendations’ – http://www.disabilitysafe.com.au/hazards-risks/manual-handling

Meittunen, E., K. .

OSHA (2009) Guidelines for Nursing Homes: Ergonomics for the Prevention of Musculoskeletal Disorders Washington: Occupational Safety and Health Administration, U.S. Department of Labor – http://www.osha.gov/ergonomics/guidelines/nursinghome/final_nh_guidelines.pdf

O'Shea, L. and M. Hennessy (2009) Transferring People Safely: Handling Clients, Residents and Clients in Health, Aged Care, Rehabilitation and DisabilityServices 3rd edition. Melbourne: WorkSafe Victoria – http://www.worksafe.vic.gov.au/wps/wcm/connect/c4c1b6804071f3a094c5dee1fb554c40/Transferring+People+Safely+-+Web.pdf?MOD=AJPERES

PAC Inc (2009a) 'Operational procedure 4.06: lifts and transfers'. Ontario: Personal Attendant Care Inc – http://www.pacdurham.ca/content/SignedHealthandSafetyOperationalProcdures/LiftsandTransfersOperationalProcn8aefts and tand tan3 c 0.0.0bd (c 0.0.0bd o>e6(40/

Two care workers in a community setting

WorkSafe Victoria (2009) Transferring People Safely: Handling Clients, Residents and Clients in Health,Aged Care, Rehabilitation and Disability Services – www.worksafe.vic.gov.au/wps/wcm/connect/c4c1b6804071f3a094c5dee1fb554c40/Transferring+People+Safely+-+Web.pdf?MOD=AJPERES

WorkSafeBC (2006a) Handle with Care: Client Handling and the Application of Ergonomics (MSI) Requirementswww2.worksafebc.com/pdfs/healthcare/HWC/HWC_L.pdf

Whittington, A. and Burns, J. (2005), 'The dilemmas of residential care staff working with the challenging behaviour of people with learning disabilities', British Journal of Clinical Psychology, 44(1), 59-76.