

Diversity of cyanobacterial biomarker genes from the stromatolites of Shark Bay, Western Australia

Tamsyn J. Garby,¹ Malcolm R. Walter,¹
Anthony W.D. Larkum² and Brett A. Neilan^{1*}

¹ School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, Georgia, USA
² School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, Georgia, USA
*Corresponding author: bneilan@gaia.com

Introduction

Summary

Families of closely related chemical compounds, which are relatively resistant to degradation, are often used as biomarkers to help trace the evolutionary history of early groups of organisms and the environments in which they lived. Biomarkers derived from hopanoid variations are particularly useful in determining bacterial community compositions. 2-Methylhopanoids have been thought to be diagnostic for cyanobacteria, and 2-methylhopanes in the geological record are taken as evidence for the presence of cyanobacteria-containing communities at the time of sediment deposition. Recently, however, doubt has been cast on the validity of 2-methylhopanes as cyanobacterial biomarkers, since non-cyanobacterial species have been shown to produce significant amounts of 2-methylhopanoids. This study examines the diversity of *hpnP*, the hopanoid biosynthesis gene coding for the enzyme that methylates hopanoids at the C2 position. Genomic DNA isolated from stromatolite-associated pustular and smooth microbial mat samples from Shark Bay, Western Australia, was analysed for bacterial diversity, and used to construct an *hpnP* clone library. A total of 117 partial *hpnP* clones were sequenced, representing 12 operational taxonomic units (OTUs). Phylogenetic analysis showed that 11 of these OTUs, representing 115 sequences, cluster within the cyanobacterial clade. We conclude that the dominant types of microorganisms with the detected capability of producing 2-methylhopanoids within pustular and smooth microbial mats in Shark Bay are cyanobacteria.

kl 1 1 kl 1 1 1 1 n kl m 1 1 kl 1 n 1 1 kl

C • rki n • i i•c ki nn ki i i•ki rki
 •ki n nn •ki / i n n • fki n i
 i n i •ki c rki n i f n rki C • w i n c
 ki i•c n ki n i rki i n • i•ki a i n -
 • n i n 11 i i • n n • n c
 rki ki nn i•c i ki C • w i n x-n
 i i • n n c rki ki nn ki i
 i x, n n f f i i % n n -
 n c n c n x n x- w i n n
 • n c rki i ki i x, n n f
 10 i i % n n c n c n x n i n
 i c n c c rki n i•ki rki -
 n c i rki n n n i i•c
 ki nn ki i i•ki w c c n n
 i•ki n n n ki nn ki i i•ki i•c
 n n n i ki i•ki
 i i n n f n i c n n n i
 i•c i n n n i n n n n n
 rki n n n i•ki x, i•ki n f n n i
 rki n n ki nn ki i i x, n n n -
 i ki i i x (f.). i•c i n i
 ki i•ki i f i•c n n c x i n
 ki nn ki i i•ki n n n f
 x i n i i f n n n c n n -
 i ki i•ki n n n f ki nn ki i•ki c
 i f i•c n i n n n x i n i i 1 p
 f n n n (n n) (n n n n n 1 . . 1 ki n n . 1 0 C 0. (•) •)- (•

с 1990 г. по 1995 г. включительно
на территории Республики Беларусь
в соответствии с Законом Республики Беларусь
от 1995 г. № 11-З «О государственном архиве»
и Положением о государственном архиве
Республики Беларусь от 1995 г. № 11-З
и Положением о государственном архиве
Республики Беларусь от 1995 г. № 11-З




Fig. 5. A world map illustrating a network of connections between various geographical locations. The map shows a dense network of connections, particularly in the North Atlantic and Europe, with significant activity also visible in the Pacific and Indian Ocean regions.

The network structure is characterized by a high density of connections, particularly in the North Atlantic and Europe. The connections are represented by colored lines and dots, indicating the strength and nature of the relationships between different geographical locations. The map shows a complex web of interactions, with significant activity also visible in the Pacific and Indian Ocean regions.

The network structure is characterized by a high density of connections, particularly in the North Atlantic and Europe. The connections are represented by colored lines and dots, indicating the strength and nature of the relationships between different geographical locations. The map shows a complex web of interactions, with significant activity also visible in the Pacific and Indian Ocean regions.

kl' m' P n' 10
(011) kl' n' x' c
n' kl' c' n' 27: 1. 00.
kl' n' P kl' n' D. (00)
x' n' n' kl' x' n' n' n' kl' x
n' n' c' kl' n' c' n' n' m' n'
273: 1.
x' C' (011) n' c' kl' n-
kl' 10' w' 1' n' D' n' n' f' n' n'
n' n' n' kl' 59: 1' 11.
w' n' P' w' (00) kl' 1' 1' 10'
1' 1' 1' f' n' kl' n' kl' 1' 1' x' n' n' x'
kl' w' kl' kl' n' 52: 0.
(1. . .) n' 1' - 10' x' n' n' f' 1' n'
1' f' kl' n' c' n' 10' n' x' n' f' kl' n' n' n' /
n' 41: . . .
n' . . . kl' 1' x' . . . n' . . . n' . . . n'
(00) n' kl' 1' 10'
1' C' n' n' n' C' n' n' kl' n' n' kl' n' f' kl' n-
1' kl' 1' kl' kl' n' 10' n' m' n' n' kl' -
n' n' n' 1' n' 1' 2: 1' . . .
n' x' w' n' c' P' n' . . . 10' n' P' (00)

kl, n, c, x, n, 169:

f, P., x, w, c, i, (00)
73:

(1, 1) kl, C, n, x, n, 173: 0

kl, n, c, x, n, (00), n, n, c, 191: